

STRONG CONVERGENCE OF A PROXIMAL POINT ALGORITHM WITH BOUNDED ERROR SEQUENCE

O. A. BOIKANYO AND G. MOROȘANU

ABSTRACT. Given any maximal monotone operator $A : D(A) \subset H \rightarrow 2^H$ in a real Hilbert space H with $A^{-1}(0) \neq \emptyset$, it is shown that the sequence of proximal iterates $x_{n+1} = (I + \gamma_n A)^{-1}(\lambda_n u + (1 - \lambda_n)x_n + e_n)$ converges strongly to the metric projection of u on $A^{-1}(0)$ for (e_n) bounded, $\lambda_n \in (0, 1)$ with $\lambda_n \rightarrow 1$ and $\gamma_n > 0$ with $\gamma_n \rightarrow \infty$ as $n \rightarrow \infty$. In comparison with our previous paper [Optim. Lett. 4 (2010), 635-641], where the error sequence was supposed to converge to zero, here we consider the classical condition that errors be bounded. In the case when A is the subdifferential of a proper convex lower semicontinuous function $\varphi : H \rightarrow (-\infty, +\infty]$, the algorithm can be used to approximate the minimizer of φ which is nearest to u .

1. INTRODUCTION

Let H be a real Hilbert space with scalar product $\langle \cdot, \cdot \rangle$ and induced norm $\| \cdot \|$. Consider the following set valued problem:

$$(1) \quad \text{find an } x \in D(A) \text{ such that } 0 \in A(x),$$

where $A : D(A) \subset H \rightarrow 2^H$ is a maximal monotone operator. We remind the reader that an operator $A : D(A) \subset H \rightarrow 2^H$ is called monotone if its graph $G(A) = \{(x, y) \in H \times H : x \in D(A), y \in Ax\}$ is a monotone subset of $H \times H$, that is, it satisfies the monotonicity property

$$\langle x - x', y - y' \rangle \geq 0, \quad \forall (x, y), (x', y') \in G(A).$$

If in addition to being monotone, the graph of A is not properly contained in the graph of any other monotone operator, then A is said to be maximal monotone. An interesting result concerning such operators is due to Minty [10] and states that a monotone operator A is maximal if and only if $(I + tA)$ is a surjection for every $t > 0$. This characterization allows one to define a single valued and nonexpansive mapping $J_t : H \rightarrow H$ by the rule $x \mapsto (I + tA)^{-1}x$ for all $t > 0$, see, e.g., [9]. The map J_t is called the resolvent of A .

One of the most effective iterative methods for solving problem (1) is the proximal point algorithm (PPA) which was initiated by Martinet [7] and further developed by Rockafellar [11]. The PPA according to Rockafellar generates a sequence (x_n) via the rule

$$(2) \quad x_{n+1} = J_{\gamma_n}(x_n + e_n), \quad \text{for all } n \geq 0$$

where $x_0 \in H$ is a given starting point, $(\gamma_n) \subset (0, \infty)$ and (e_n) is a sequence of computational errors. Since the PPA converges only weakly in general, to a solution of problem (1), see [1, 4], modifying the PPA in order to generate strongly convergent sequences has

2000 *Mathematics Subject Classification.* 47J25, 47H05, 47H09.

Key words and phrases. maximal monotone operator, proximal point algorithm, nonexpansive map, resolvent operator, prox-Tikhonov method.

been the subject of intense research. Recently, Lehdili and Moudafi [6] combined the proximal method (2) with the Tikhonov regularization to obtain the iterative process

$$(3) \quad x_{n+1} = J_{\gamma_n}(\alpha_n x_n + e_n), \quad \text{for all } n \geq 0$$

where $x_0 \in H$ is a given starting point, $\gamma_n \in (0, \infty)$ and $\alpha_n \in (0, 1)$ with the requirement $\alpha_n \rightarrow 1$ as $n \rightarrow \infty$, and (e_n) is a sequence of computational errors. Under the summability condition on $(\|e_n\|)$ and some additional conditions on α_n and γ_n , they showed that the sequence generated by (3) converges strongly to a point of $A^{-1}(0)$ nearest to u provided that this set is not empty. Xu [13] extended the prox-Tikhonov method (3) to

$$(4) \quad x_{n+1} = J_{\gamma_n}(\lambda_n u + (1 - \lambda_n)x_n + e_n), \quad \text{for all } n \geq 0$$

for given $x_0, u \in H$, $\gamma_n \in (0, \infty)$, $\lambda_n \in (0, 1)$, and (e_n) has the same meaning as above. Worth mentioning is the fact that when $\lambda_n \rightarrow 0$ and $\|e_n\| \rightarrow 0$ as $n \rightarrow \infty$, algorithm (4) above is equivalent to an iterative process of the form

$$(5) \quad x_{n+1} = \lambda_n u + (1 - \lambda_n)J_{\gamma_n}x_n + e_n, \quad \text{for all } n \geq 0,$$

see [2] for details. The scheme (5) was independently shown by Xu [12] and Kamimura and Takahashi [5] to be strongly convergent to a point of $A^{-1}(0)$ (if this set is not empty) which is nearest to u , provided that $\lambda_n \rightarrow 0$ with $\sum_{n=0}^{\infty} \lambda_n = \infty$, $(\|e_n\|)$ is summable and $\gamma_n \rightarrow \infty$. In [2], the authors extended this result to general errors which only converge to zero in norm.

In this paper, we pursue a different direction to derive a strong convergence result associated with algorithm (4). More precisely, we will show that if $A^{-1}(0) \neq \emptyset$, $\lambda_n \rightarrow 1$, (e_n) is bounded and $\gamma_n \rightarrow \infty$, then the sequence generated by (4) converges strongly to an element of $A^{-1}(0)$ which is nearest to u .

2. MAIN RESULT

Our analysis relies on the following lemma which describes the asymptotic behavior of the resolvent of a maximal monotone operator. It was proved independently by Bruck Jr. [3] and Moroșanu [8].

Lemma 1. *Let $A : D(A) \subset H \rightarrow 2^H$ be a maximal monotone operator with $\emptyset \neq F := A^{-1}(0)$. Then for any $u \in H$, $(I + tA)^{-1}u \rightarrow P_F u$ as $t \rightarrow \infty$, where $P_F u$ denotes the projection of u on F .*

We will also need the following lemma in proving our main result

Lemma 2. *Assume that $A : D(A) \subset H \rightarrow 2^H$ is a maximal monotone operator with $F := A^{-1}(0) \neq \emptyset$. For any fixed $x_0, u \in H$, let (x_n) be generated by*

$$(6) \quad x_{n+1} = J_{\gamma_n}(\lambda_n u + (1 - \lambda_n)(x_n + e_n)), \quad \text{for all } n \geq 0,$$

where $\lambda_n \in (0, 1)$ and $\gamma_n \in (0, \infty)$. If $(1 - \lambda_n)\|e_n\|/\lambda_n \leq C$ for some $C > 0$, then (x_n) is bounded.

Proof. Denote $w_n := \lambda_n u + (1 - \lambda_n)(x_n + e_n)$. Then for each $p \in A^{-1}(0)$, we have from (6) and the fact that the resolvent operator is nonexpansive

$$\begin{aligned} \|x_{n+1} - p\| &= \|J_{\gamma_n} w_n - J_{\gamma_n} p\| \\ &\leq \|w_n - p\|. \end{aligned}$$

Therefore, it would suffice to show that the sequence (w_n) is bounded. Note that

$$(7) \quad w_{n+1} = \lambda_{n+1} u + (1 - \lambda_{n+1})e_{n+1} + (1 - \lambda_{n+1})J_{\gamma_n} w_n, \quad \text{for all } n \geq 0.$$

Let $M > 0$ be big enough such that for some $p \in F$

$$\|u - p\| + \frac{(1 - \lambda_n)}{\lambda_n} \|e_n\| \leq M \quad \text{and} \quad \|w_0 - p\| \leq 2M, \quad \text{for all } n \geq 0.$$

Then from (7), we see that

$$\begin{aligned} \|w_{n+1} - p\|^2 &= \|\lambda_{n+1}(u - p) + (1 - \lambda_{n+1})e_{n+1} + (1 - \lambda_{n+1})(J_{\gamma_n}w_n - p)\|^2 \\ &\leq (1 - \lambda_{n+1})^2 \|J_{\gamma_n}w_n - p\|^2 \\ &\quad + 2\lambda_{n+1} \left\langle u - p + \frac{(1 - \lambda_{n+1})}{\lambda_{n+1}} e_{n+1}, w_{n+1} - p \right\rangle \\ (8) \quad &\leq (1 - \lambda_{n+1})^2 \|w_n - p\|^2 + 2M\lambda_{n+1} \|w_{n+1} - p\|, \end{aligned}$$

where the first inequality follows from the subdifferential inequality

$$\|x + y\|^2 \leq \|y\|^2 + 2\langle x, x + y \rangle \quad \text{for all } x, y \in H$$

and the last inequality follows from the fact that the resolvent operator is nonexpansive. Now, if we assume that

$$(9) \quad \|w_n - p\| \leq 2M$$

for some $n \in \mathbb{N}_0$, then from (8), we have

$$\begin{aligned} (\|w_{n+1} - p\| - \lambda_{n+1}M)^2 &\leq 4(1 - \lambda_{n+1})^2 M^2 + \lambda_{n+1}^2 M^2 \\ &\leq (2(1 - \lambda_{n+1})M + \lambda_{n+1}M)^2 \\ &= (2M - \lambda_{n+1}M)^2, \end{aligned}$$

showing that (9) also holds true for $n+1$. Therefore, we conclude that (w_n) is bounded. \square

We are now in a position to prove our main result

Theorem 3. *Assume that $A : D(A) \subset H \rightarrow 2^H$ is a maximal monotone operator with $F := A^{-1}(0) \neq \emptyset$. For any fixed $x_0, u \in H$, let the sequence (x_n) be generated by (6) where $\lambda_n \in (0, 1)$ and $\gamma_n \in (0, \infty)$ for all $n \geq 0$. If (e_n) is bounded, $\lambda_n \rightarrow 1$ and $\gamma_n \rightarrow \infty$ as $n \rightarrow \infty$, then (x_n) converges strongly to $P_F u$, the metric projection of u on F .*

Proof. We know from Lemma 2 that (x_n) is bounded. Moreover, from (6), we have

$$\begin{aligned} \|x_{n+1} - P_F u\| &\leq \|x_{n+1} - J_{\gamma_n} u\| + \|J_{\gamma_n} u - P_F u\| \\ &\leq (1 - \lambda_n) \|x_n - u + e_n\| + \|J_{\gamma_n} u - P_F u\| \end{aligned}$$

where the second inequality follows from the fact that the resolvent operator is nonexpansive. The result follows immediately on passing to the limit in the above inequality. \square

Remark 4. We point out that for (e_n) bounded and $\lambda_n \rightarrow 1$ as $n \rightarrow \infty$, algorithms (6) and (7) are not equivalent. Note that under these assumptions and $A^{-1}(0) \neq \emptyset$, Lemma 2 guarantees that the sequence (w_n) defined above is bounded. Consequently, for any $u \notin A^{-1}(0) =: F$, (w_n) converges strongly to $u \neq P_F u$, whereas (x_n) generated from (6) converges strongly to $P_F u$ (see Theorem 3 above).

Remark 5. In [2], it was proved that if $F := A^{-1}(0) \neq \emptyset$, then the sequence generated by algorithm (6) converges strongly to $P_F u$, provided that $\lambda_n \rightarrow 0$ with $\sum_{n=0}^{\infty} \lambda_n = \infty$, $\gamma_n \rightarrow \infty$ and $\|e_n\|/\lambda_n \rightarrow 0$. We note that since $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$ in [2, Theorem 1], for very large n , the argument of the resolvent operator in (6) becomes arbitrarily close to that of algorithm (2). In this paper, we have employed a different approach: instead of the assumption $\lambda_n \rightarrow 0$, we have used the condition $\lambda_n \rightarrow 1$ as $n \rightarrow \infty$ which allows

the argument of the resolvent operator in (6) to tend to the point u as $n \rightarrow \infty$. This approach is based on the fact that for very large values of t , $(I + tA)^{-1}$ approximates the zero of A which is closest to u . Theorem 3 is a strong theoretical result, under the classic boundedness condition on errors. It remains to see whether computationally, the speed/rate of convergence of algorithm (6) is fast enough.

REFERENCES

- [1] H. H. Bauschke, J. V. Burke, F. R. Deutsch, H. S. Hundal and J. D. Vanderwerff, *A new proximal point iteration that converges weakly but not in norm*, Proc. Amer. Math. Soc. 133 (2005), no. 6, 1829-1835.
- [2] O. A. Boikanyo and G. Moroşanu, *A proximal point algorithm converging strongly for general errors*, Optim. Lett. 4 (2010), 635-641.
- [3] R. E. Bruck Jr., *A strongly convergent iterative solution of $0 \in U(x)$ for a maximal monotone operator U in Hilbert space*, J. Math. Anal. Appl. 48 (1974), 114-126.
- [4] O. Güler, *On the convergence of the proximal point algorithm for convex minimization*, SIAM J. Control Optim. 29 (1991), 403-419.
- [5] S. Kamimura and W. Takahashi, *Approximating solutions of maximal monotone operators in Hilbert spaces*, J. Approx. Theory 106 (2000), 226-240.
- [6] N. Lehdili and A. Moudafi, *Combining the proximal algorithm and Tikhonov regularization*, Optimization 37 (1996), 239-252.
- [7] B. Martinet, *Régularisation d'inéquations variationnelles par approximations successives*, Rev. Française Informat. Recherche Opérationnelle 4 (1970), Ser. R-3, 154-158.
- [8] G. Moroşanu, *Asymptotic behaviour of resolvent for a monotone set in a Hilbert space*, Atti Accad. Nsz. Lincei 61 (1977), 565-570.
- [9] ———, *Nonlinear Evolution Equations and Applications*, Reidel, Dordrecht, 1988.
- [10] G. J. Minty, *Monotone (nonlinear) operators in Hilbert space*, Duke Math. J. 29 (1962), 341-348.
- [11] R. T. Rockafellar, *Monotone operators and the proximal point algorithm*, SIAM J. Control Optim. 14 (1976), 877-898.
- [12] H. K. Xu, *Iterative algorithms for nonlinear operators*, J. London Math. Soc. (2) 66 (2002), 240-256.
- [13] ———, *A regularization method for the proximal point algorithm*, J. Glob. Optim. 36 (2006), 115-125.

DEPARTMENT OF MATHEMATICS AND ITS APPLICATIONS, CENTRAL EUROPEAN UNIVERSITY, NADOR
U. 9, H-1051 BUDAPEST, HUNGARY

E-mail address: boikanyo_oganeditse@ceu-budapest.edu, and morosanug@ceu.hu